

The prevalence of post-operative Seroma formation after Ventral Hernia Repair in patients using Abdominal Binder versus not using Abdominal-Binder

Shahidullah Ahmad, Sheikh Muhammad Ibqar Azeem, Kaleemullah, Waleed Mabood, Sahibzada Salma Rahman, Usman Ullah

Abstract:

Background: Ventral hernias are a commonly encountered pathology in surgical units. These are abnormal projection of abdominal viscera or part of viscera through a defect in anterior abdominal wall. Seroma formation is one of the common post operative complications. Wide spread dissection and the resultant enlarged dead space results in the collection of serous fluid formation. The objective of this study was to compare the prevalence of post-operative seroma formation in patients after ventral hernia repair with use of abdominal binder versus no abdominal binder.

Materials and Methods: This randomized controlled trail was conducted in the department of general surgery, Mercy Teaching Hospital Peshawar, Pakistan from January 2021 to December 2022. 108 patients were randomly allocated by toss method into 2 groups i.e., abdominal belt binding group and non-binding group having 54 patients in each group. The data for the sample were described by counts and percentages and for the population as confidence interval with 95% confidence level. Mc Nemar chi-square test was used to assess post-operative seroma formation after ventral hernia repair in abdominal binder using versus non-using abdominal binder

Results: Seroma formation occurred in 27% (15) ($15 \times 100 / 54 = 27$) patients in abdominal binder group, while in non-binder group seroma formation occurred in 75% (41) ($41 \times 100 / 54 = 75$) patients and H_{01} was accepted as results were statistically not significant ($p\text{-value} > 0.05$).

Conclusion: In our study it was found that post-operative seroma formation was reduced in patients using abdominal binders as compared to non-binders after ventral hernia repair. So, this study will help surgeons in alleviating post-operative seroma formation by using abdominal binder post-operatively.

Keywords: Seroma, ventral hernia, epigastric hernia, abdominal binder, complications.

Introduction:

Ventral hernias are a commonly encountered pathology in surgical units.¹ These are abnormal projection of abdominal viscera or part of viscera through a defective anterior abdominal wall.² The prevalence of ventral hernias is increasing rapidly, mostly due to obesity and requires surgical management in elderly patients.³

The options for ventral hernia repair are dependent on the size of defect, patient's priority and

surgeons' preference.³ Guidelines published by the IEHS implies open hernia repair for small defects < 2cm and open mesh repair is devised for defects >10cm while the rest of hernias are considered for laparoscopic approach.⁴ Laparoscopic hernia repair is considered sophisticated in patients with large hernias or having adhesions from previous abdominal operations.⁵ However both techniques of surgical repair are considered secure and are widely employed for ventral hernia repair.⁶⁻⁷

Received

date: 14th July, 2023

Accepted

date: 2nd Novemer, 2023

Peshawar Medical
College, Peshawar

S Ahmad
SMI Azeem
W Mabood
SS Rahman

DHQ, Hospital Dagar,
Buner

Kaleemullah

Correspondence:

Dr. Sheikh Muhammad
Ibqar Azeem
Associate Professor
Surgery, Department of
Surgery, Peshawar Medical
College, Peshawar.
Cell No: +00 000-0000000
email: ibqarazeem@gmail.
com

Mesh support decreases the recurrence rate in ventral hernias.⁸ Successful hernia repair implies no post-operative complications i.e., pain, seroma formation, wound infection, scar formation and recurrence.⁹

Seroma formation is one of the common post operative complications.¹⁰ Wide spread dissection and the resultant enlarged dead space results in the collection of fluid and seroma formation.¹¹ Clinically profound seroma is found in about 3-17% of laparoscopic ventral hernia repair cases.¹²⁻¹³ Seromas can persist for several months or even years and can cause post-operative infection, skin flap necrosis, delayed hospital discharge and recurrence as well.¹⁴⁻¹⁶

In case of asymptomatic seroma “wait and watch” policy is employed, however if seroma enlarges or show signs of infection then aspiration or even re-exploration becomes necessary.¹⁷

Zia et al.¹⁸ from Daru Sehat Hospital Karachi, Pakistan for the period of 1 and half years from Jan 2017 to June 2019 found that seroma formation occurred in 4(8%) cases in abdominal binder group and in 12(24%) cases in non-binder group & the results were statistically significant (p-value<0.05).

Dogar et al.¹⁹ from Central Park Teaching Hospital Lahore, Pakistan for the period of 6 months from January 2019 to June 2019 found that seroma formation occurred in 2(6.7%) patients in abdominal binder group and 8(26.7%) patients in non-binder group and the results were statistically significant (p-value<0.05).

Unawareness about the outcome after abdominal binder use and no abdominal binder on post-operative seroma formation after ventral hernia repair was our research problem. Unavailable data regarding this problem was our knowledge gap and what is the outcome after abdominal binder use and no abdominal binder on post-operative seroma formation after ventral hernia repair.

The objective of this study was to compare the prevalence of post-operative seroma formation

after ventral hernia repair in abdominal binder using versus not using abdominal binder.

Research Null Hypothesis (H_{01}): H_{01} there is no statistically significant difference in seroma formation in ventral hernia repair patients post-operatively in abdominal binder using versus not using abdominal binder.

Material and Methods:

Design, setting and duration: This randomized controlled trail was conducted in the department of general surgery, Mercy Teaching Hospital Peshawar, Pakistan for a period of 2 years from Jan 2021 to December 2022. Approval was granted from Hospital ethical committee and informed consent was taken from patients or attendants.

Sampling and Randomization: All adult patients having primary or recurrent umbilical and para-umbilical hernia and planned to undergo open ventral hernia repair under general anesthesia were included in the study. Patients having chronic liver disease, Ascites, Renal Failure, Pregnant females or ventral hernia with gut perforation or strangulation were excluded from the study. 108 patients were randomly allocated by toss method into 2 groups i.e., abdominal belt binding group and non-binding group having 54 patients in each group.

Procedure, intervention and follow-up: All 108 patients under went ventral hernia repair by open ventral hernia repair technique. Pre-operatively patients were given IV antibiotics injection Ceftriaxone 1gm, 1 hour before incision. All patients were given Injection Ceftriaxone 1gm IV* BD for 1 day post-operatively and then oral Antibiotic Tab Moxifloxacin 400mg* OD for 5 days. All patients were given Injection Toradol* BD for 1 day post-operatively and then Tab Voren 50mg* BD for pain relief for 5 days. Patients were discharged after 24 hours of surgery. 54 patients in Abdominal binder group were applied abdominal binder immediate post-operative period 24 hourly for 1 week then >12 hours for 6 weeks and in non-binder group no abdominal belt was applied.

Table 1: Comparison of seroma formation after ventral hernia repair in abdominal binder using versus not using of abdominal binder

Groups		Seroma Formation			Chi-square value	d.f	p-value
		Yes	No	Total			
Abdominal Binder Use	Yes	15	39	54	108	1	0.851
	No	41	13	54			

H01 accepted at α 0.05

Seroma formation was measured on 7th post-operative day using trans-abdominal ultrasound in both groups.

Data collection and analysis plan: Gender (men/women) and age groups (≤ 40 years and ≥ 40 years) were our matching variables, whereas compare the outcome after abdominal binder use and no abdominal binder on post-operative seroma formation after ventral hernia repair was our research variable. The data for the sample were described by counts and percentages and for the population as confidence interval with 95% confidence level. Mc Nemar chi-square test was used to assess post-operative seroma formation after ventral hernia repair in abdominal binder using versus non-using abdominal binder. Data was analyzed by SPSS v.17 (SPSS Inc, Chicago, IL, USA).

Results:

In our study the mean age in abdominal binder group was 41.7 ± 7.5 years, while it was 43.5 ± 6.5 years in non-binder group. There were 29 women and 25 men in abdominal binder group, whereas 21 women and 33 men in Non-Binder group. There were 30 Para-umbilical hernia cases and 24 Epigastric hernia cases in abdominal binder group while 32 Para-umbilical hernia cases and 22 Epigastric hernia cases in Non-Binder group.

Seroma formation occurred in 27% (15) ($15 \times 100 / 54 = 27$) patients in abdominal binder group, while in non-binder group seroma formation occurred in 75% (41) ($41 \times 100 / 54 = 75$) patients and H01 was accepted as results were statistically not significant (p -value > 0.05). (Table 1)

Discussion:

Post-operative seroma formation occurs after a number of abdominal surgeries e.g., Exploratory laparotomy, Cesarean section, Ventral hernia repairs & abdominoplasty etc. Morales-Conde et al.²⁰ found that the incidence of seroma formation after ventral hernia repair is 46%. Lund et al.²¹ retrieved seroma formation in 22% patient post laparoscopic ventral hernia repair.

A number of methods have been devised to decrease the incidence of post-operative seroma formation. One of the methods to reduce the chances of seroma formation is sclerotherapy, which constitute filling of seroma cavity with a sclerosant that induces fibrosis and seals the dead space.²² Other methods to prevent seroma formation are negative pressure drains, sealing agents e.g., collagen sponge having coating of human coagulation factors and pressure dressing or abdominal binder use. External pressure reduces fluid leak from vessels, so, abdominal binder can help reduce seroma formation.¹¹

Abdominal binders are tightening belts that encompass the abdomen to enhance recovery process especially after exploratory laparotomy, Cesarean section, bariatric surgeries, hysterectomy or ventral hernia repair surgeries.²³ Abdominal binder usage is safe after abdominal surgeries and they minimally effect respiratory dynamics and intra-abdominal pressure.²⁴

In our study it was found that seroma formation occurred in 27% patients in abdominal binder group, while in non-binder group seroma formation occurred in 75% patients. The results were not statistically significant.

Similar to our study, Zia et al.¹⁸ from Daru Sehat Hospital Karachi, Pakistan found that seroma formation occurred in 4(8%) cases in abdominal binder group and in 12(24%) cases in non-binder group and the results were statistically significant (p -value < 0.05).

Likewise Dogar et al.¹⁹ from Central Park Teaching Hospital Lahore, Pakistan found that seroma formation occurred in 2(6.7%) patients in ab-

dominal binder group and 8(26.7%) patients in non-binder group & the results were statistically significant (p-value<0.05).

Conclusion:

In our study it was found that post-operative seroma formation was reduced in patients using abdominal binders as compared to non-binders. So, this study will help surgeons in alleviating post-operative seroma formation by using abdominal binder post-operatively. Further it will prevent complications associated with seroma formation i.e., infections, necrosis and revision surgeries.

Conflict of interest: None

Funding source: None

Role and contribution of authors:

Shahidullah Ahmad, conception and acquisition of data, critical revision and final approval.

Sheikh Muhammad Ibqar Azeem, conception and acquisition of data, drafting the manuscript, literature search and final approval.

Kaleemullah, conception and acquisition of data, critical revision.

Waleed Mabood, literature search and data analysis.

Sahibzada Salma Rahman, drafting the manuscript and literature search.

Usman Ullah, collected the references and also helped in material and methods writing.

References:

- Bell-Allen N, Swift K, Sontag NJ, O'Rourke N. Ventral hernia repair with a hybrid laparoscopic technique. *ANZ J Surg* 2022 Oct;92(10):2529-33.
- Hospital Episode Statistics. UK ventral hernia data: <https://digi-tal.nhs.uk/data-and-information/publications/statistical/hospital-admitted-patient-care-activity/2017-18>.
- Shelton J, Poulouse BK, Phillips S, Moore D, Nealon W, Penson D et al. Epidemiology and cost of ventral hernia repair: making the case for hernia research. *Hernia* 2012;16:179-183.
- Bittner R, Bingener-Casey J, Dietz U, Fabian M, Ferzli GS, Fortelny RH et al. Guidelines for laparoscopic treatment of ventral and incisional abdominal wall hernias (international Endo hernia society [IHES]). *Surg Endosc* 2014; 28: 353-79.
- Zhang Y, Zhou H, Chai Y, Cao C, Jin K, Hu Z. Laparoscopic versus open incisional and ventral hernia repair: a systematic

- review and meta-analysis. *World J. Surg* 2014; 38: 2233-40.
- Al Chalabi H, Larkin J, Mehigan B, McCormick P. A systematic review of laparoscopic versus open abdominal incisional hernia repair, with meta-analysis of randomized controlled trials. *Int J Surg* 2015; 20: 65-74.
- Heniford BT, Park A, Ramshaw BJ, Voeller G. Laparoscopic repair of ventral hernias: nine years' experience with 850 consecutive hernias. *Ann Surg* 2003; 238: 391-9.
- Poulouse BK (2018) So what's the chance of this mesh causing me a problem in the long run? *Ann Surg* 267: e66. <https://doi.org/10.1097/SLA.0000000000002681>
- Sandø A, Rosen MJ, Heniford BT, Bisgaard T. Long-term patient-reported outcomes and quality of the evidence in ventral hernia mesh repair: a systematic review. *Hernia* 2020 Aug;24:695-705.
- Shermak MA, Rotellini-Coltvet LA, Chang D. Seroma development following body contouring surgery for massive weight loss: patient risk factors and treatment strategies. *Plast Reconstr Surg* 2008 July1; 122(1): 280-288.
- Moshe D, Karen IG. Preventing Postoperative Seroma formation in Abdominal wall Hernia by Intraoperative Hypertonic Saline Irrigation, early Report. *Open Access J Surg* 2019;10(2).
- Lund S, Farley D. A decade of experience with laparoscopic ventral hernia repairs. *The Am J Surg* 2019 Mar 1;217(3):546-9.
- Clemens MW, Butler CE. Ventral abdominal hernia repair: technique—external oblique release. *The SAGES Manual of Hernia Surgery* 2019:217-36.
- He C, Lu J, Ong MW, Lee DJ, Tan KY, Chia CL. Seroma prevention strategies in laparoscopic ventral hernia repair: a systematic review. *Hernia*. 2020 Aug;24:717-31.
- Kumar S, Lal B, Misra MC. Post-mastectomy seroma: a new look into the etiology of an old problem. *JR Coll Surg Edinb* 1995 Oct 1;40(5):292-4.
- Sood A, Kotamarti VS, Therattil PJ, Lee ES. Sclerotherapy for the management of seromas: a systematic review. *Eplasty* 2017;17.
- Schachtrupp A, Klinge U, Junge K, Rosch R, Bhardwaj RS, Schumpelick V. Individual inflammatory response of human blood monocytes to mesh biomaterials. *Br J Surg* 2003 Jan; 90(1):114-120. doi:10.1007/s00268-005-7970-2
- ZIA MK, SEHRISH Y, ABBASI F, BANO S, NEHAL Z, QURESHI A. Comparison of Abdominal Binder Use Versus No Abdominal Binder Use After Ventral Abdominal Wall Hernia Repair.
- Dogar MA, Chaudhery WM, Rasheed G, Butt AS, Riaz A, Ayaz M. comparison of abdominal binder use versus no abdominal binder use after ventral abdominal wall hernia repair. A randomized controlled trial. *PPMJ* 2018;29(3):111-5.
- Morales Conde S, Suarez Artacho G, Socas Macias M, Baranco Moreno A. Retro prosthetic seroma after laparoscopic ventral hernia repair: incidence, risk factors and clinical significance. *Hernia* 2015 Dec;19(6): 943-947.
- Lund S, Farley D. A decade of experience with laparoscopic ventral hernia repairs. *The Am J Surg* 2019 Mar 1;217(3):546-9.
- Penaud A, Quignon R, Danin A, Bahé L, Zakine G. Alcohol sclerodhesis: an innovative treatment for chronic Morel-Lavallée lesions. *J Plast ReconstrAesthetSurg* 2011 Oct 1;64(10):e262-4.
- Gustafsson UO, Scott MJ, Schwenk W, Demartines N, Roulin D, Francis N et al. Guidelines for perioperative care in elective colonic surgery: Enhanced Recovery After Surgery (ERAS®) Society recommendations. *World J Surg* 2013, 37:259-284. 10.1007/s00268-012-1772-0 4
- Cheifetz O, Lucy SD, Overend TJ, Crowe J. The effect of abdominal support on functional outcomes in patients following major abdominal surgery: a randomized controlled trial. *Physiother Can* 2010, 62:242-253. 10.3138/physio.62.3.242