


Aligning Care Delivery with Evidence-Based Standards: Bridging the Gaps

Integrating evidence-based practices into healthcare delivery systems remains pivotal in achieving optimal patient outcomes. Evidence-Based Medicine (EBM) integrates clinical expertise, patient values, and the best available evidence to inform decision-making processes for individual patient care.¹ However, the alignment of care delivery with these principles often encounters systemic challenges, necessitating deliberate efforts to ensure adherence to standardized protocols and guidelines.




Hierarchy of Evidence:²

The foundation of evidence-based practice lies in the hierarchy of evidence, where higher levels of study designs, such as randomized controlled trials and systematic reviews, provide more rigorous and reliable results with minimal bias. As healthcare continues to evolve, leveraging these robust sources becomes integral to informed decision-making.

Defining Standard Care Delivery:^{3,4}

Standard care delivery is characterized by the

consistent provision of high-quality healthcare services based on established protocols, guidelines, and patient safety standards. Its key objectives include improving patient outcomes, ensuring safety, optimizing resource utilization, and fostering patient satisfaction. The core pillars—access, quality, safety, efficiency, and patient-centered care—form the bedrock of a resilient healthcare system.



Components of Standard Care Delivery:^{5,6}

1. Clinical Standards: Evidence-based protocols for diagnosis and treatment.
2. Operational Standards: Streamlined workflow processes, staffing, and resource management.
3. Patient Experience: Emphasis on effective communication, empathy, and care coordination.
4. Technology Integration: Utilization of Electronic Medical Records (EMR) and Health Information Management Systems (HIMS).
5. Regulatory Compliance: Adherence to national and international healthcare standards

Importance of Standard Care Delivery:^{7,8}

Adherence to standard care delivery ensures equitable access, minimizes errors, and fosters trust between patients and providers. Furthermore, it aligns with accreditation requirements, such as those set forth by the Joint Commission International and Safe Care standards, enhancing efficiency and reducing costs.

Examples of Standard Practices:^{9,10}

- Timely and accurate patient assessments.
- Safe medication management adhering to the “5 Rights of Medication.”
- Infection prevention protocols, including hand hygiene and sterilization.
- Consistent documentation and reporting systems.
- Regular audits and quality control measures.

Challenges and Breaches in Care Delivery:^{11,12}

Despite established standards, breaches often occur due to insufficient staff training, outdated technology, communication failures, and resource shortages. Common breaches include delayed care, medication errors, infection control lapses, and incomplete documentation. These incidents lead to increased morbidity, legal repercussions, and diminished trust.

Strategies for Improvement:^{4,13}

To mitigate these challenges, healthcare systems must:

- Prioritize regular staff training and education.
- Implement advanced EMR and HIMS systems.
- Strengthen quality control and auditing practices.
- Cultivate a culture of safety and accountability.
- Maintain continuous monitoring and feedback mechanisms.

Conclusion and Call to Action:

Ensuring alignment with evidence-based standards is imperative for achieving quality, safety, and efficiency in healthcare. As healthcare professionals, we must identify gaps in current processes, leverage technology-driven solutions, and foster a culture of teamwork and continuous learning. Together, we can bridge the gaps and deliver the highest standard of care.

Prof Syed Asim Ali Jaffary

Editor

FRCS (Edinburgh), Master in Hepatobiliary Surgery (France), Fellowship in Liver Transplantation (Korea), Diploma in Medical Education (Dundee), EMBA Healthcare Management (DUHS)

Consultant Surgeon and

Associate Chief Medical Officer

Aga Khan University, Karachi

asim.jaffary@aku.edu

Ph: +92 (0)3002401292

References:

1. Sackett DL, Rosenberg WM, Gray JA, Haynes RB, Richardson WS. Evidence based medicine: what it is and what it isn't. *BMJ*. 1996;312(7023):71-72.
2. Centre for Evidence-Based Medicine. Levels of Evidence (March 2009). University of Oxford. Available from: <https://www.cebm.ox.ac.uk/resources/levels-of-evidence/levels-of-evidence-march-2009>.
3. World Health Organization. Quality of Care: A Process for Making Strategic Choices in Health Systems. Geneva: WHO; 2006.
4. Institute of Medicine. Crossing the Quality Chasm: A New Health System for the 21st Century. Washington, DC: National Academies Press; 2001.
5. Donabedian A. The quality of care: How can it be assessed? *JAMA*. 1988;260(12):1743-1748.
6. National Patient Safety Foundation. Free from Harm: Accelerating Patient Safety Improvement Fifteen Years after To Err Is Human. Boston: NPSF; 2015.
7. Berwick DM, Nolan TW, Whittington J. The Triple Aim: care, health, and cost. *Health Aff (Millwood)*. 2008;27(3):759-769.
8. World Health Organization. Patient Safety: Making Health Care Safer. Geneva: WHO; 2017.
9. Joint Commission International. Joint Commission International Accreditation Standards for Hospitals. 7th ed. Oak Brook, IL: Joint Commission Resources; 2021.
10. Centers for Disease Control and Prevention (CDC). Guide to Infection Prevention for Outpatient Settings: Minimum Expectations for Safe Care. Atlanta: CDC; 2016.
11. Reason J. Human error: models and management. *BMJ*. 2000;320(7237):768-770.
12. World Health Organization. Global Patient Safety Action Plan 2021-2030. Geneva: WHO; 2021.
13. National Academy of Medicine. Vital Directions for Health and Health Care: Priorities from a National Academy of Medicine Initiative. Washington, DC: NAM; 2015.